

Acoustic analogs of two-dimensional black holes

Mariano Cadoni*

*Dipartimento di Fisica, Università di Cagliari, and INFN sezione di Cagliari,
Cittadella Universitaria 09042 Monserrato, ITALY*

We present a general method for constructing acoustic analogs of the black hole solutions of two-dimensional (2D) dilaton gravity. Because by dimensional reduction every spherically symmetric, four-dimensional (4D) black hole admits a 2D description, the method can be also used to construct analogue models of 4D black holes. We also show that after fixing the gauge degrees of freedom the 2D gravitational dynamics is equivalent to an one-dimensional fluid dynamics. This enables us to find a natural definition of mass M , temperature T and entropy S of the acoustic black hole. In particular the first principle of thermodynamics $dM = TdS$ becomes a consequence of the fluid dynamics equations. We also discuss the general solutions of the fluid dynamics and two particular cases, the 2D Anti-de sitter black hole and the 4D Schwarzschild black hole.

I. INTRODUCTION

In recent years it has become increasingly clear that condensed matter systems can be used to mimic various kinematical aspects of general relativity [1, 2, 3, 4, 5]. Analogue models of gravity have been used not only to mimic black holes and event horizons but also to describe other spacetime structures, cosmological solutions and field theory in curved spacetimes [6, 7, 8, 9, 10, 11, 12, 13, 14]. Condensed matter analogs of gravity could be also used to shed light on old and new puzzling features of the gravitational interaction such as the information loss problem in the Hawking evaporation of a black hole or the holographic principle. Moreover, analogue models of gravity that use condensed matter systems have also the nice feature of being, at least in principle, experimentally testable in laboratory. In the near future this could open the way to a “black hole phenomenology” based not on astrophysical but condensed matter experiments.

One strong limitation of this approach is that until now analogue models of black holes can be used to mimic the kinematical but not the dynamical aspects of gravitational systems. For instance, one can use acoustic analogs to study the kinematics of event horizons, or even the Hawking radiation but not the formation of horizons due to some distribution of matter or the back-reaction of the geometry on the Hawking radiation (some progress along this direction has been achieved in Ref. [11, 12]). A further unpleasant consequence of this limitation is the impossibility to reproduce black hole thermodynamics using analogue models of black holes. It is well known that the laws of black hole thermodynamics are a consequence of the Einstein equations. Although one can define the temperature associated with an acoustic horizon [3], it is extremely problematic to find the right definition of mass and entropy of an acoustic black hole.

At first sight to mimic the dynamical aspects of general relativity using a condensed matter system seems an impossible task. The description of the gravitational system is characterized by a huge redundancy of gauge degrees of freedom and by a separation between matter sources and gravitational field (respectively described by the stress-energy tensor and the curvature tensors in the Einstein equations). Conversely, focusing on acoustic black holes, the fluid dynamics is characterized by few physical degrees of freedom (pressure, velocity, density) and does not seem to allow for a source-field description.

Nonetheless, there is a situation in which one could hope to merge gravitational and fluid dynamics. A spherically symmetric, black hole with gauge degrees of freedom completely fixed is characterized, owing to Birkhoff’s theorem, by a handful of physical observables (the charges associated with global symmetries). These observables could be put in correspondence with the fluid parameters. A spherically symmetric black hole can be described by an effective two-dimensional(2D) dilaton gravity model [15] obtained from four-dimensional(4D) Einstein gravity by retaining only the radial modes. Thus, the most promising framework for trying to merge gravitational and fluid dynamics is that of 2D gravity.

In this paper we will show that this is possible. We propose a general method for constructing acoustic analogs of the black hole solutions of two-dimensional (2D) dilaton gravity (Section II). Further, we show that after fixing the gauge degrees of freedom the 2D gravitational dynamics is equivalent to a constrained one-dimensional fluid dynamics

*Electronic address: mariano.cadoni@ca.infn.it

(Section III). This enables us to find a natural definition of mass M , temperature T and entropy S of the acoustic black hole. In particular the first principle of thermodynamics $dM = TdS$ becomes a consequence of the fluid dynamics equations (Sections IV and V). The constraint for the fluid dynamics can be enforced in two independent ways. If the external parameters for the fluid (profile of the flux tube or potential for the external forces) are given the constraint simply determines the barotropic equation of state for the fluid. Alternatively, if one chooses a given Equation of state the constraint determines the form of the external parameters for the fluid (Section VI). We also discuss two particular cases, the 2D Anti-de sitter black hole and the 4D Schwarzschild black hole (Section VII).

II. 2D ACOUSTIC METRIC AND 2D BLACK HOLES

Let us consider the generic 2D dilaton gravity model (For a review see Ref. [15]), characterized by a dilaton potential V

$$A = \frac{1}{2} \int d^2x \sqrt{-g} (\phi R + \lambda^2 V(\phi)), \quad (1)$$

where ϕ is a scalar (the dilaton) and λ is parameter with dimensions of a length⁻¹ (throughout this paper we will use natural units $c = \hbar = k_B = 1$). Two-dimensional dilaton gravity models have been used in various situations as effective description of 4D gravity. In particular, every spherically symmetric 4D solution can be described in terms of a 2D dilaton gravity model. The model (1) admits black hole solutions, which in the Schwarzschild gauge take the form [16],

$$ds^2 = - \left(J(\lambda r) - \frac{2M}{\lambda} \right) d\tau^2 + \left(J(\lambda r) - \frac{2M}{\lambda} \right)^{-1} dr^2, \quad \phi = \lambda r, \quad (2)$$

where M is the black hole mass and $J = \int V d\phi$. The black hole horizon is located at $r = r_h$, with $J(\lambda r_h) = 2M/\lambda$.

We want to find acoustic analogs of the 2D black hole solutions (2). Obviously, the most natural candidate is a fluid, whose motion is essentially one-dimensional. Let us therefore consider the steady, locally irrotational flow of a 3D fluid which is barotropic and inviscid. If the transverse velocities (in the z and y directions) are small with respect to the velocity along the x axis, we can set them to zero. The propagation of acoustic disturbances χ on the background fields is determined, at the linearized level, by the wave equation $\nabla^2 \chi = (1/\sqrt{-g}) \partial_\mu (\sqrt{-g} g^{\mu\nu} \partial_\nu \chi) = 0$, where $g_{\mu\nu}$ is the so-called acoustic metric [3]. In the case under consideration, the acoustic metric describes a one-dimensional slab geometry where the velocity is always along the x direction and the velocity profile depends only on x ,

$$ds^2 = \frac{\bar{\rho}_0}{c} [- (c^2 - v_0^2) dt^2 - 2v_0 dx dt + dx^2], \quad (3)$$

where $\bar{\rho}_0(x), v_0(x)$ are, respectively, the background density and velocity of the fluid and $c(x) = \sqrt{(dP/d\bar{\rho}_0)}$ is the local speed of sound (P is the pressure).

The two metrics (2), (3) can be transformed one into the other using a Painlevé Gullstrand-like coordinate transformation [17, 18] and identifying the function J and the black hole mass M in terms of the physical parameters of the fluid. It is convenient to work with a dimensionless fluid density. This can be achieved by multiplying the metric (3) with a factor λ^{-4} and defining the dimensionless density $\rho_0 = \lambda^{-4} \bar{\rho}_0$. The coordinate transformations and the identification relating (2) with (3) are,

$$\begin{aligned} r &= \int \rho_0 dx, \\ \tau &= t + \int dx \frac{v_0}{c^2 - v_0^2}, \\ J &= \frac{2M}{\lambda} + \frac{\rho_0}{c} (c^2 - v_0^2). \end{aligned} \quad (4)$$

Notice that the coordinate r (or equivalently the dilaton ϕ) from the point of view of the fluid plays the role of an extensive quantity. If the flux tube has a constant section it is proportional to the mass of the fluid contained between some reference initial point and point x . We stress the fact that the correspondence between 2D black holes (2) and the acoustic metric (3) is exact. This is to be compared with what happens for 4D static black holes, for which the correspondence with the acoustic metric can be established only up to a conformal transformation of the metric [3].

Eqs. (4) allow to associate a one-dimensional slab geometry (and a corresponding acoustic black hole) to every 2D black hole solution of the dilaton gravity model (1). They can be also used to construct acoustic analogs of 4D spherically symmetric black hole solutions (e.g Schwarzschild). This is not surprising, being the causal structure of a 4D spherically symmetric black hole completely encoded in its (r, t) sections.

III. 2D GRAVITATIONAL DYNAMICS VERSUS FLUID DYNAMICS

In the previous section we have seen that given a generic 2D black hole we can always construct a 2D acoustic analog. However, the correspondence is purely kinematical. In order to formulate the correspondence at a dynamical level we need to compare the gravitational dynamics with the dynamics of the fluid, which determines ρ_0, v_0, c .

At the dynamical level, the most striking difference between the two physical systems is that the gravitational dynamics has to be covariant under general coordinate transformations. In particular, this means that we have a huge redundancy of gauge degrees of freedom in the description of our 2D gravity system. These gauge degrees of freedom have to be fixed if we want to construct a dynamical acoustic analogue of the black hole. On the other hand, 2D dilaton gravity is purely topological. We do not have any physical propagating gravitational degree of freedom. This leaves open the possibility that once the gauge freedom have been fixed, the dynamics for the resulting global gravitational degrees of freedom could match fluid dynamics.

The field equations for the 2D dilaton gravity model (1) are

$$R = -\lambda^2 \frac{dV}{d\phi}, \quad \nabla_\mu \nabla_\nu \phi - \frac{1}{2} g_{\mu\nu} \lambda^2 V = 0. \quad (5)$$

We consider only static solutions and fix the diffeomorphisms invariance, choosing the Schwarzschild gauge for the 2D metric

$$ds^2 = -X(r)d\tau^2 + X^{-1}(r)dr^2. \quad (6)$$

Using Eq. (6) into the field equations (5), one finds after some manipulations that the field equation for the dilaton ϕ are given simply by $d\phi/dr = \lambda$, which implies that the dilaton is $\phi = \lambda r$. The field equations for the global gravitational degree of freedom $X(r)$ become instead

$$\frac{dX}{dr} = \lambda V. \quad (7)$$

One can also easily check that the black hole (2) is solution of Eq. (7).

The fundamental equations describing fluid dynamics are the Euler and the continuity equations. The steady flow of the fluid giving rise to the one-dimensional slab geometry described in the previous section, is therefore governed by the equations:

$$\bar{\rho}_0 v_0 \frac{dv_0}{dx} + \frac{dP}{dx} + \bar{\rho}_0 \frac{d\psi}{dx} = 0, \quad \bar{\rho}_0(x) v_0(x) A(x) = \text{const.}, \quad (8)$$

where P is the pressure, ψ denotes the potential for external driving forces acting on the fluid (including Newtonian gravity) and $A(x)$ is the area of the section of the flux tube. We will consider separately the two cases: a) The external potential ψ is non-homogeneous and the section of the flux tube is constant. b) The external potential is zero and the section of the flux tube does depend on x .

Let us first focus on case a). Using the barotropicity condition $P = P(\rho_0)$, passing from the coordinate x to the coordinate r as defined in Eq. (4) and defining the new variables

$$\begin{aligned} X &= \frac{\rho_0}{c} (c^2 - v_0^2), \\ Y &= \rho_0 c, \\ F &= \ln \left(\frac{c}{\rho_0} \right), \end{aligned} \quad (9)$$

the Euler and continuity equations (8) take, respectively, the form

$$\begin{aligned} \frac{dX}{dr} &= 2 \frac{dY}{dr} - X \frac{dF}{dr} + 2e^{-F} \frac{d\psi}{dr} \\ \frac{d}{dr} [Y(Y - X)] &= 0. \end{aligned} \quad (10)$$

One can now easily realize that the fluid dynamics equations (10) can be made equivalent to the gravitational equations (7) just by introducing the constraint

$$2 \frac{dY}{dr} - X \frac{dF}{dr} + 2e^{-F} \frac{d\psi}{dr} = \lambda V(\phi). \quad (11)$$

In fact, when constraint (11) is enforced and the variable X is identified in terms of the gravitational variables as in Eq. (6) the Euler equation (10) matches exactly Eq. (7).

Simple counting of the degrees of freedom and of the fundamental equations of the dynamics reveals that we can consistently map the fluid dynamics into the gravitational dynamics imposing constraints. After gauge fixing, the gravitational system has only one degree of freedom, the function X parametrizing the metric in Eq. (6) and one equation. The scalar field ϕ is constrained to be proportional to r and therefore represents just the spacelike coordinate of the 2D spacetime. On the other hand, if the external potential is given, the fluid dynamics is characterized by three unknown functions c, ρ_0, v_0 and two equations, so that we have the freedom of imposing an additional constraint. If the external potential ψ has been fixed, imposing the constraint (11) is implicitly equivalent to an equation of state $P = P(\rho_0)$ for the fluid. From this point of view, the equation of state for the fluid may be seen as a consequence of the gravitational dynamics. This is very similar to Jacobson's interpretation of the Einstein equation as an equation of state [19]. Alternatively, if one first chooses a given equation of state the constraint (11) becomes an equation that determines the external potential ψ .

Two-dimensional gravity allows us to define the mass M of the black hole solution (2) as a scalar, which on-shell becomes constant [20]

$$M = \frac{1}{2\lambda} \left(\lambda^2 \int V(\phi) d\phi - (\nabla\phi)^2 \right). \quad (12)$$

Owing to the correspondence between the dynamics of 2D dilaton gravity and the previously discussed constrained fluid dynamics, we also expect the existence of a function Γ of the fluid parameters X, Y, F, ψ which is constant (independent of the flux tube coordinate x) by virtue of Eqs. (10) and (11). This function is given by

$$\Gamma = a \left[\int_{r_0}^r (-X dF + 2e^{-F} d\psi) + 2Y - X \right], \quad (13)$$

where a and r_0 are arbitrary constants. One can easily check that $d\Gamma/dr = 0$ follows from Eq. (10) and (11).

Eq. (13) allows us to find the acoustic counterpart of the black hole mass M . Fixing the constant $a = \lambda/2$, using Eq. (11) and reading from Eq. (4) the solution for X , $X = J - 2M/\lambda$, one readily finds $M = \Gamma$.

IV. BLACK HOLE THERMODYNAMICS AND FLUID DYNAMICS

It is a well known fact that black hole thermodynamics follows directly from the gravitational field equations, once one has defined appropriately the thermodynamical parameters. In the case of 2D gravity the thermodynamical state of the black hole is completely defined by temperature T , energy (mass) M and entropy S . The mass is given by Eq. (12). The temperature can be defined in terms of the periodicity of the euclidean metric at the horizon, whereas the entropy can be computed using Noether charge techniques [21] and turns out to be proportional to the dilaton evaluated at the black hole horizon. We have

$$T = \frac{\lambda}{4\pi} V(\phi_h), \quad S = 2\pi\phi_h. \quad (14)$$

The first principle of thermodynamics follows directly from Eq. (7). In fact, using Eqs. (2) one has $S = 2\pi J^{-1}(2M/\lambda)$, which in turn implies $dM = (\lambda/4\pi)(dJ(r_h)/d\phi_h)dS$. On the other hand Eq. (7) implies $dJ/d\phi = V$, which inserted in the previous equation and using Eq. (14) gives $dM = TdS$.

The correspondence between gravitational and fluid dynamics described in the previous section allows us to define straightforwardly the thermodynamical parameters, T_a, M_a, S_a associated with the acoustic black hole. Moreover, the first principle of thermodynamics will become consequence of the fluid dynamics equations (10),(11).

Although the phenomenon of Hawking radiation for acoustic black holes is a rather involved issue [22], one can still define also for the acoustic black hole a temperature in terms of the ‘‘surface gravity’’ or the periodicity of the Euclidean section at the acoustic horizon $c = v_0$. We have

$$T_a = \frac{1}{4\pi} \left(\frac{dX}{dr} \right)_{c=v_0} = \frac{1}{4\pi} \left[\frac{1}{c} \frac{d}{dx} (c^2 - v_0^2) \right]_{c=v_0}. \quad (15)$$

The above defined temperature coincides with the usual definition of temperature for acoustic black holes [3]. Owing to the constraint (11), in our 2D case we can also write down an other, completely equivalent, expression for the temperature,

$$T_a = \left[\frac{1}{4\pi\rho_0} \left(2\frac{dY}{dx} - X\frac{dF}{dx} + 2e^{-F}\frac{d\psi}{dx} \right) \right]_{c=v_0}. \quad (16)$$

For generic acoustic black holes it is not so easy to find a natural definition of mass. In the case under consideration we have a natural candidate for M_a , given by the quantity (13), which remains constant along the flux tube. Because it is a constant we can as well define M_a as Eq. (13) evaluated at the acoustic horizon ,

$$M_a = \frac{\lambda}{2} \left[\int_{r_0}^{r_h} dr \left(-X \frac{dF}{dr} + 2e^{-F} \frac{d\psi}{dr} \right) + 2Y(r_h) \right], \quad (17)$$

where r_0 is chosen such that $M_a = M$. The entropy of the acoustic black hole can be defined by analogy with the gravitational black hole, $S_a = 2\pi\phi_h = 2\pi r_h$. Using the coordinate transformation (4) we get,

$$S_a = 2\pi \int_{r=\infty}^{r_h} dx \rho_0(x), \quad (18)$$

where we have fixed the arbitrary integration constant in such way that the integration domain is the whole range of the coordinate x outside the acoustic horizon. The entropy for the acoustic black hole is proportional to the total mass of the fluid outside the acoustic horizon. Notice that this definition corresponds to the intuitive notion of entropy as an extensive quantity, which counts elementary degrees of freedom.

One can easily show, differentiating Eq. (17) and using Eqs. (16) and (18) that the thermodynamical parameters satisfy the first principle $dM_a = T_a dS_a$.

We can also show that the first principle can be derived directly from our constrained fluid dynamics. Evaluating the constraint (11) at the acoustic horizon ($r = r_h$) we get $\lambda dJ(r_h) = (2dY/dr - X dF/dr + 2e^{-F} d\psi/dr)_{r_h} dr_h$. Using Eqs. (16) and (18) together with $dJ(r_h) = (2/\lambda)dM$, we find $dM_a = T_a dS_a$.

V. FLUX TUBE WITH NON-CONSTANT SECTION

Until now we have considered a flux tube of constant section with an non-homogeneous external potential. The discussion can be easily extended to a flux tube with non-constant section. In this latter case the presence of a external potential is an unnecessary complication. For this reason we only consider the dynamics of a fluid with zero external potential and a non-constant section. In this case the fluid dynamics equations (8), written in terms of the variables defined in Eq. (9) are

$$\frac{dX}{dr} = 2 \frac{dY}{dr} - X \frac{dF}{dr}, \quad \frac{d}{dr} [(Y - X) Y A^2] = 0. \quad (19)$$

Introducing the constraint

$$2 \frac{dY}{dr} - X \frac{dF}{dr} = \lambda V, \quad (20)$$

Eqs. (19) can be made equivalent to the gravitational equations (7).

The function that is constant owing to Eqs. (19), (20) and can be therefore interpreted as the black hole mass is now

$$\gamma = a \left[\int_{r_0}^r (-X dF) + 2Y - X \right]. \quad (21)$$

Also in this case, once $A(x)$ has been fixed, the constraint (20) is equivalent to an equation of state for the fluid. On the other hand if the equation of state of the fluid is given, Eq. (20) determines $A(x)$. In this latter case, Eq. (20) represents a “geometrical constraint” for the fluid flow of the same kind of that discussed in Ref. [6] in relation to the realizability of acoustic horizons.

The mass and temperature of the acoustic black hole are now

$$M_a = \frac{\lambda}{2} \left[\int_{r_0}^{r_h} dr \left(-X \frac{dF}{dr} \right) + 2Y(r_h) \right], \quad T_a = \left[\frac{1}{4\pi\rho_0} \left(2 \frac{dY}{dx} - X \frac{dF}{dx} \right) \right]_{c=v_0}, \quad (22)$$

whereas the entropy is always given by Eq. (18). One can easily check the validity of the first principle of thermodynamics and that this principle is a consequence of the equations of the fluid dynamics (19), (20).

VI. SOLUTIONS OF THE CONSTRAINED FLUID DYNAMICS

Apart from its intrinsic theoretical interest, our approach could also have a direct experimental realization. Given a four-dimensional, spherically symmetric, gravitational black hole, one can derive by dimensional reduction the corresponding 2D dilaton gravity model (1) and related 2D black hole (2). The 2D black hole solution is completely characterized by the function J and the mass M . Once $J(r)$ and M are given, we can construct explicitly the corresponding 2D acoustic black hole and, at least in principle, realize the experimental set-up, which will produce the acoustic black hole.

In order to have a well-defined physical system, we need to solve the fluid dynamics equations, i.e we need to know ρ_0, c, v_0 as function of the flux tube coordinate x . Equations (7) and the second equation in (10) (or in (19)) can be easily integrated

$$X = J - \frac{2M}{\lambda}, \quad Y = \frac{1}{2} \left(X + \sqrt{X^2 + \alpha^2} \right), \quad (23)$$

where α is a constant in case a), whereas $\alpha = \text{const}/A(x)$ in case b).

To solve the dynamics completely, the constraint (11) (or 20) has to be enforced. This can be done in two nonequivalent ways. 1. One can choose a given barotropic equation of state for the fluid and the constraint (11) ((20)) determines the external potential ψ (the profile $A(x)$ for the flux tube). 2. The external potential ψ (the flux tube profile $A(x)$) is given and the constraint (11) ((20)) determines the barotropic equation of state for the fluid. Because one usually operates with a fluid with a given equation of state, the physically relevant situation is obviously the first one. Therefore we will consider here only possibility 1.. To handle with a definite and common situation, we will consider a fluid with a constant speed of sound c . The barotropic equation of state is $P = c^2 \rho_0$, which expressed in terms of the variables (9) gives

$$Y e^F = c^2. \quad (24)$$

Let us first consider the case of flux tube of constant section (case a)). Inserting Eqs. (24) and (23) into the constraint (11) and using Eqs. (9),(23), (24) one obtains the solution for $v_0(X), \rho_0(X)$ and $\psi(X)$,

$$v_0 = \frac{\alpha c}{2Y}, \quad \rho_0 = \frac{1}{c} Y, \quad \psi(X) = c^2 \left(\frac{X}{2Y} - \ln Y \right), \quad (25)$$

where $Y(X)$ is given as in Eq. (23). The acoustic horizon is located at $X = 0$ ($Y = \alpha/2$). At the horizon the density remains finite, $\rho_0 = \alpha/2c$. This is the best we can do in the general case. The calculations can be completed once a specific 2D gravitational model (a specific function J) is given. Taking into account that $\rho_0 = dr/dx$, the previous equation for ρ_0 becomes a differential equation, which determines the coordinate transformation $r = r(x)$.

Let us now consider the case of a flux tube on non-constant section (our case b)). We can use Eqs. (23) to obtain

$$A^2 \propto \frac{1}{Y(Y-X)}. \quad (26)$$

Notice that $Y \geq X$. In order to determine $Y(X)$ we use Eq. (24) in the constraint (20) to get

$$2 \frac{dY}{dX} + \frac{X}{Y} \frac{dY}{dX} = 1. \quad (27)$$

The solution of this equation is given in implicit form by

$$X = 2Y \ln Y, \quad (28)$$

where for simplicity we have set to zero the integration constant. We can now write the solution of our fluid dynamics equation in the form

$$v_0 = c \sqrt{1 - \frac{X}{Y}}, \quad \rho_0 = \frac{Y}{c}, \quad (29)$$

where $Y(X)$ is implicitly defined by Eq. (28). The horizon is located at $X = 0$, the subsonic region for $X > 0$. From very general arguments (see for instance Ref. [5]) one expects that the acoustic horizon forms at a minimum of the

area of the flux tube section (the so called Laval nozzle). Let us show that this is also true in our case. From Eqs. (26) and (27) one finds

$$A' \propto \frac{(1 - 2Y')}{Y - X}, \quad (30)$$

where the prime denotes derivation with respect to X . On the other hand, Eq. (27) tells us that at the horizon we have $Y' = 1/2$ and that in the subsonic (supersonic) region $Y' < 1/2$ ($Y' > 1/2$). It follows immediately that $A' = 0$ at the horizon and that $A' > 0$ ($A' < 0$) in the subsonic (supersonic) region, i.e that in the horizon region the flux tube must take the form of a Laval nozzle.

VII. TWO EXAMPLES: 2D ANTI-DE SITTER AND 4D SCHWARZSCHILD BLACK HOLE

In this section we will consider two particular cases as illustration of our general procedure . Our first example is the 2D anti-de Sitter (AdS) black hole [23]. The corresponding 2D dilaton gravity model (1) has a constant potential $V(\phi) = 2$, from which follows $J(r) = \lambda^2 r^2$. The black hole solution (2) describes a 2D AdS spacetime with an horizon at $r = \sqrt{2M/\lambda^3}$. The corresponding acoustic black hole can be easily constructed using our general procedure. Let us consider separately the two cases of constant and non-constant flux tube section.

A. Flux tube with constant section

Considering a flux tube in presence of a non-homogeneous external potential and with constant section, the acoustic analog of the 2D AdS black hole can be easily found inserting $X = \lambda^2 r^2 - 2M/\lambda$ into Eqs. (25). Using the expression for ρ_0 given in equation (25) one can also find the relation between the radial coordinate r of the gravitational black hole and the spatial coordinate x of the acoustic black hole. The $0 \leq r < r_h$ interior of the AdS black hole is described by the supersonic region of the acoustic black hole. At $r = 0$ the fluid parameters ρ_0, v_0 and the external potential are finite. For $r > 0$ v_0 decreases and ρ_0 grows monotonically. Also the external potential ψ grows for $r > 0$ but reaches its maximum at the horizon $r = r_h$. At the acoustic horizon, $v_0 = c$ and both the density ρ_0 and the external potential remain finite. The asymptotic region $r \rightarrow \infty$ of the gravitational black hole (the timelike conformal boundary of the 2D AdS spacetime) corresponds to the subsonic region of the acoustic black hole. Going in this region $v_0 \rightarrow 0$, the fluid density ρ_0 and fluid pressure P diverge as expected owing to the continuity equation. Conversely, the potential ψ of the external force acting on the fluid diverges logarithmically as $-\ln r$. The behavior of the parameters ψ, v_0, ρ_0 , normalized to their horizon values and for $\alpha = 1$ are shown in Fig. (1) as a function of the variable $X = \lambda^2 r^2 - 2M/\lambda$.




FIG. 1: Behavior of the normalized parameters ψ/c^2 (normal line), $v_0/v_0(r_h)$ (bold line) and $\rho_0/\rho_0(r_h)$ (dashed line) as a function of the variable X for a flux tube with constant section and non-homogeneous external potential characterized by $\alpha = 1$. The velocity and fluid density are normalized to their horizon values. The acoustic horizon is located at $X = 0$

B. Flux tube with non-constant section

If we use a flux tube of non-constant section to mimic the 2D AdS black hole, we get the typical shape of a converging/diverging Laval nozzle. We can find the solution for the fluid equations using $X = \lambda^2 r^2 - 2M/\lambda$ into

Eqs. (26), (28) (29). In this case the fluid can be used to describe only the horizon region of the AdS black hole, but not the full spacetime. In fact, if A^2 of Eq. (26) has to stay positive X and because of Eq. (28) it follows that $-2/e \leq X \leq \sqrt{e}$ and correspondingly $1/e \leq Y \leq \sqrt{e}$. The supersonic region of the acoustic black hole describes the region $\sqrt{2M/\lambda^3 - 2/(e\lambda^2)} = r_m \leq r < r_h = \sqrt{2M/\lambda^3}$ of the AdS black hole interior. Analogously, the subsonic region can be used to describe only the region $\sqrt{2M/\lambda^3} = r_h < r \leq r_M = \sqrt{2M/\lambda^3 + \sqrt{e}/\lambda^2}$ of the AdS black hole.

The profile of the flux tube is that of the Laval nozzle depicted in Fig. (2), which also shows the behavior of the fluid velocity and fluid density, normalized to their horizon values, as a function of the variable Y defined in Eq. (28). The flux tube section A starts at a finite value at $r = r_m$ decreases in the supersonic region of the acoustic black hole till it reaches its minimum $A(r_h)$ at the horizon. In the subsonic region $r_h < r < r_M$, A grows and diverges at $r = r_M$. The fluid velocity decreases monotonically starting from $v_0(r_m) = \sqrt{3}c$ in the supersonic region to reach first its horizon value $v_0(r_h) = c$ and then $v_0(r_M) = 0$, as expected owing to the equation of continuity. Finally, the fluid density ρ_0 , grows monotonically from $\rho_0(r_m) = (ec)^{-1}$ in the supersonic region to the horizon value $\rho_0(r_h) = c^{-1}$ to reach $\rho_0(r_M) = \sqrt{e}c^{-1}$ in the subsonic region.




FIG. 2: Behavior of the normalized parameters $A/A(r_h)$ (normal line), $v_0/v_0(r_h)$ (bold line) and $\rho_0/\rho_0(r_h)$ (dashed line) as a function of the variable Y for a flux tube with non-constant section. The flux tube section, the velocity and fluid density are normalized to their horizon values. The acoustic horizon is located at $Y = 1$

Our second example is the 4D Schwarzschild black hole. By means of a dimensional reduction, retaining only the radial modes, 4D Einstein gravity can be described by the 2D gravity model (1) with dilaton potential $V(\phi) = 1/\sqrt{2\phi}$ and with λ^{-2} identified as the 4D Newton constant G . The 2D black hole solution (2) becomes,

$$ds^2 = \left(\sqrt{2\lambda r} - \frac{2M}{\lambda} \right) d\tau^2 + \left(\sqrt{2\lambda r} - \frac{2M}{\lambda} \right)^{-1} dr^2. \quad (31)$$

This solution describes the 2D sections of the 4D Schwarzschild solution $ds^2 = -(1 - 2MG/R)d\tau^2 + (1 - 2MG/R)^{-1}dR^2$. In fact the metric (31) can be put in the Schwarzschild form by means of the Weyl rescaling of the 2D metric $g_{\mu\nu} \rightarrow (1/\sqrt{2\phi})g_{\mu\nu}$ and the coordinate transformation $r = (\lambda/2)R^2$. Mass, temperature and entropy of the 2D black hole are invariant under Weyl transformations of the metric [24]. In fact the temperature and entropy of the 2D black hole (31), given by $T = \lambda^2/8\pi M$, $S = (4\pi M^2/\lambda^2)$, match exactly those of the 4D Schwarzschild black hole after setting $\lambda^{-2} = G$.

The acoustic analogue of the Schwarzschild black hole can be constructed, working with a flux tube of constant section putting $X = \sqrt{2\lambda r} - 2M/\lambda = \lambda R - 2M/\lambda$ into Eqs. (25), (or into Eqs. (26), (28), (29) if working with a flux tube of non constant section).

The features of the acoustic Schwarzschild black hole are qualitatively similar to those of the previously discussed AdS acoustic black hole.

VIII. SUMMARY AND OUTLOOK

In this paper we have developed a general method for constructing acoustic analogs of 2D black hole solutions. The procedure can be also used for 4D black holes that admit an effective 2D description. In particular this is the case of spherically symmetric 4D black holes. We have also shown that after fixing the gauge degrees of freedom, the 2D gravitational dynamics is equivalent to a constrained fluid dynamics. The correspondence between gravitational and fluid dynamics has allowed a natural definition of the thermodynamical parameters mass, temperature and entropy

of the acoustic black hole. Moreover, we have shown that the first principle of thermodynamics follows from the constrained fluid dynamics.

The results presented in this paper represent just a first step for finding condensed matter systems that mimic gravitational dynamics. It is not hard to identify the two main (and related) limitations of our approach. First, our discussion holds for static gravitational configurations. In many interesting situations, for instance the computation of the back-reaction for an evaporating black hole, one has to deal with non static solutions. The generalization of our method to the non static case may be extremely non trivial, in particular the fixing of the gravitational gauge degrees of freedom necessary to reproduce the fluid dynamics. The simplicity of 2D approach can be very useful also for treating the non static case. Second, in this paper we have considered a 2D gravitational model without matter. The absence of sources for the gravitational field represent a strong simplification for the dynamical equations. On the other hand, we need the presence of sources if we want to describe a realistic situation. Finding a fluid dynamic counterpart of the matter stress-energy tensor appearing in the gravitational field equations may not be a easy task.

From a more technical point of view, if we work with a fluid with a given equation of state the practical realization of a physical system that mimics the 2D black hole requires a particular careful choice of an external potential (flux tube of constant section) or of the flux tube profile (flux tube of nonconstant section).

Acknowledgments

I am grateful to A. Fabbri, S. Liberati a S. Mignemi for helpful comments.

-
- [1] W. G. Unruh, Phys. Rev. Lett. **46** (1981) 1351.
 - [2] M. Visser, arXiv:gr-qc/9311028.
 - [3] M. Visser, Class. Quant. Grav. **15** (1998) 1767 [arXiv:gr-qc/9712010].
 - [4] M. Novello, M. Visser and G. Volovik, “Artificial Black Holes” (World Scientific, 2002)
 - [5] M. Visser, C. Barcelo and S. Liberati, Gen. Rel. Grav. **34** (2002) 1719 [arXiv:gr-qc/0111111].
 - [6] C. Barcelo, S. Liberati and M. Visser, Int. J. Mod. Phys. A **18** (2003) 3735 [arXiv:gr-qc/0110036].
 - [7] P. O. Fedichev, and U. R. Fischer, Phys. Rev. A **69** (2004) 033602 [arXiv:cond-mat/0303063].
 - [8] P. O. Fedichev, and U. R. Fischer, Phys. Rev. Lett. **91** (2003) 240407 [arXiv:cond-mat/0304342].
 - [9] C. Barcelo, S. Liberati and M. Visser, Int. J. Mod. Phys. D **12** (2003) 1641 [arXiv:gr-qc/0305061].
 - [10] C. Barcelo, S. Liberati, S. Sonogo and M. Visser, arXiv:gr-qc/0408022.
 - [11] R. Balbinot, S. Fagnocchi, A. Fabbri and G. P. Procopio, arXiv:gr-qc/0405096.
 - [12] R. Balbinot, S. Fagnocchi and A. Fabbri, arXiv:gr-qc/0405098.
 - [13] E. Berti, V. Cardoso and J. P. S. Lemos, arXiv:gr-qc/0408099.
 - [14] V. Cardoso, J. P. S. Lemos and S. Yoshida, arXiv:gr-qc/0410107.
 - [15] D. Grumiller, W. Kummer and D. V. Vassilevich, Phys. Rept. **369** (2002) 327 [arXiv:hep-th/0204253].
 - [16] D. Louis-Martinez and G. Kunstatter, Phys. Rev. D **49** (1994) 5227.
 - [17] P. painlevé, C. R. Acad. Sci. (paris) **173** (1921) 677.
 - [18] A. Gullstrand, Arkiv. Mat. Astron. Fys. **16(8)** (1922) 1.
 - [19] T. Jacobson, Phys. Rev. Lett. **75** (1995) 1260 [arXiv:gr-qc/9504004].
 - [20] R. B. Mann, Phys. Rev. D **47** (1993) 4438 [arXiv:hep-th/9206044].
 - [21] R. C. Myers, Phys. Rev. D **50** (1994) 6412 [arXiv:hep-th/9405162]. [22]
 - [22] W. G. Unruh and R. Schutzhold, arXiv:gr-qc/0408009.
 - [23] M. Cadoni and S. Mignemi, Phys. Rev. D **51** (1995) 4319 [arXiv:hep-th/9410041].
 - [24] M. Cadoni, Phys. Lett. B **395** (1997) 10 [arXiv:hep-th/9610201].